


**AP[®] CALCULUS AB
2001 SCORING GUIDELINES**

Question 1

Let R and S be the regions in the first quadrant shown in the figure above. The region R is bounded by the x -axis and the graphs of $y = 2 - x^3$ and $y = \tan x$. The region S is bounded by the y -axis and the graphs of $y = 2 - x^3$ and $y = \tan x$.



- (a) Find the area of R .
 (b) Find the area of S .
 (c) Find the volume of the solid generated when S is revolved about the x -axis.

Point of intersection

$$2 - x^3 = \tan x \text{ at } (A, B) = (0.902155, 1.265751)$$

(a) Area $R = \int_0^A \tan x \, dx + \int_A^{\sqrt[3]{2}} (2 - x^3) \, dx = 0.729$

or

$$\text{Area } R = \int_0^B ((2 - y)^{1/3} - \tan^{-1} y) \, dy = 0.729$$

or

$$\text{Area } R = \int_0^{\sqrt[3]{2}} (2 - x^3) \, dx - \int_0^A (2 - x^3 - \tan x) \, dx = 0.729$$

(b) Area $S = \int_0^A (2 - x^3 - \tan x) \, dx = 1.160$ or 1.161

or

$$\text{Area } S = \int_0^B \tan^{-1} y \, dy + \int_B^2 (2 - y)^{1/3} \, dy = 1.160 \text{ or } 1.161$$

or

$$\begin{aligned} \text{Area } S &= \int_0^2 (2 - y)^{1/3} \, dy - \int_0^B ((2 - y)^{1/3} - \tan^{-1} y) \, dy \\ &= 1.160 \text{ or } 1.161 \end{aligned}$$

(c) Volume $= \pi \int_0^A ((2 - x^3)^2 - \tan^2 x) \, dx$
 $= 2.652\pi$ or 8.331 or 8.332

3 : { 1 : limits
1 : integrand
1 : answer

3 : { 1 : limits
1 : integrand
1 : answer

3 : { 1 : limits and constant
1 : integrand
1 : answer

AP[®] CALCULUS AB 2002 SCORING GUIDELINES

Question 1

Let f and g be the functions given by $f(x) = e^x$ and $g(x) = \ln x$.

- (a) Find the area of the region enclosed by the graphs of f and g between $x = \frac{1}{2}$ and $x = 1$.
- (b) Find the volume of the solid generated when the region enclosed by the graphs of f and g between $x = \frac{1}{2}$ and $x = 1$ is revolved about the line $y = 4$.
- (c) Let h be the function given by $h(x) = f(x) - g(x)$. Find the absolute minimum value of $h(x)$ on the closed interval $\frac{1}{2} \leq x \leq 1$, and find the absolute maximum value of $h(x)$ on the closed interval $\frac{1}{2} \leq x \leq 1$. Show the analysis that leads to your answers.

(a) Area = $\int_{\frac{1}{2}}^1 (e^x - \ln x) dx = 1.222$ or 1.223

2 { 1 : integral
1 : answer

(b) Volume = $\pi \int_{\frac{1}{2}}^1 ((4 - \ln x)^2 - (4 - e^x)^2) dx$
 = 7.515π or 23.609

4 { 1 : limits and constant
2 : integrand
< -1 > each error
Note: 0/2 if not of the form
 $k \int_a^b (R(x)^2 - r(x)^2) dx$
1 : answer

(c) $h'(x) = f'(x) - g'(x) = e^x - \frac{1}{x} = 0$
 $x = 0.567143$

3 { 1 : considers $h'(x) = 0$
1 : identifies critical point
and endpoints as candidates
1 : answers

Absolute minimum value and absolute maximum value occur at the critical point or at the endpoints.

$h(0.567143) = 2.330$

$h(0.5) = 2.3418$

$h(1) = 2.718$

The absolute minimum is 2.330.

The absolute maximum is 2.718.

Note: Errors in computation come off the third point.

AP[®] CALCULUS AB
2007 SCORING GUIDELINES

Question 1

Let R be the region in the first and second quadrants bounded above by the graph of $y = \frac{20}{1+x^2}$ and below by the horizontal line $y = 2$.

- (a) Find the area of R .
- (b) Find the volume of the solid generated when R is rotated about the x -axis.
- (c) The region R is the base of a solid. For this solid, the cross sections perpendicular to the x -axis are semicircles. Find the volume of this solid.

$$\frac{20}{1+x^2} = 2 \text{ when } x = \pm 3$$

(a) Area = $\int_{-3}^3 \left(\frac{20}{1+x^2} - 2 \right) dx = 37.961$ or 37.962

1 : correct limits in an integral in (a), (b), or (c)

2 : $\begin{cases} 1 : \text{integrand} \\ 1 : \text{answer} \end{cases}$

(b) Volume = $\pi \int_{-3}^3 \left(\left(\frac{20}{1+x^2} \right)^2 - 2^2 \right) dx = 1871.190$

3 : $\begin{cases} 2 : \text{integrand} \\ 1 : \text{answer} \end{cases}$


(c) Volume = $\frac{\pi}{2} \int_{-3}^3 \left(\frac{1}{2} \left(\frac{20}{1+x^2} - 2 \right) \right)^2 dx$
 $= \frac{\pi}{8} \int_{-3}^3 \left(\frac{20}{1+x^2} - 2 \right)^2 dx = 174.268$

3 : $\begin{cases} 2 : \text{integrand} \\ 1 : \text{answer} \end{cases}$

AP[®] CALCULUS AB
2012 SCORING GUIDELINES

Question 2

Let R be the region in the first quadrant bounded by the x -axis and the graphs of $y = \ln x$ and $y = 5 - x$, as shown in the figure above.



- (a) Find the area of R .
- (b) Region R is the base of a solid. For the solid, each cross section perpendicular to the x -axis is a square. Write, but do not evaluate, an expression involving one or more integrals that gives the volume of the solid.
- (c) The horizontal line $y = k$ divides R into two regions of equal area. Write, but do not solve, an equation involving one or more integrals whose solution gives the value of k .

$$\ln x = 5 - x \Rightarrow x = 3.69344$$

Therefore, the graphs of $y = \ln x$ and $y = 5 - x$ intersect in the first quadrant at the point $(A, B) = (3.69344, 1.30656)$.

(a)
$$\text{Area} = \int_0^B (5 - y - e^y) dy$$

$$= 2.986 \text{ (or } 2.985)$$

OR

$$\text{Area} = \int_1^A \ln x \, dx + \int_A^5 (5 - x) \, dx$$

$$= 2.986 \text{ (or } 2.985)$$

(b)
$$\text{Volume} = \int_1^A (\ln x)^2 \, dx + \int_A^5 (5 - x)^2 \, dx$$

(c)
$$\int_0^k (5 - y - e^y) \, dy = \frac{1}{2} \cdot 2.986 \text{ (or } \frac{1}{2} \cdot 2.985)$$

3 : $\left\{ \begin{array}{l} 1 : \text{integrand} \\ 1 : \text{limits} \\ 1 : \text{answer} \end{array} \right.$


3 : $\left\{ \begin{array}{l} 2 : \text{integrands} \\ 1 : \text{expression for total volume} \end{array} \right.$

3 : $\left\{ \begin{array}{l} 1 : \text{integrand} \\ 1 : \text{limits} \\ 1 : \text{equation} \end{array} \right.$

AP[®] CALCULUS AB
2013 SCORING GUIDELINES

Question 5

Let $f(x) = 2x^2 - 6x + 4$ and $g(x) = 4\cos\left(\frac{1}{4}\pi x\right)$. Let R be the region bounded by the graphs of f and g , as shown in the figure above.



- (a) Find the area of R .
- (b) Write, but do not evaluate, an integral expression that gives the volume of the solid generated when R is rotated about the horizontal line $y = 4$.
- (c) The region R is the base of a solid. For this solid, each cross section perpendicular to the x -axis is a square. Write, but do not evaluate, an integral expression that gives the volume of the solid.

$$\begin{aligned} \text{(a) Area} &= \int_0^2 [g(x) - f(x)] dx \\ &= \int_0^2 \left[4\cos\left(\frac{\pi}{4}x\right) - (2x^2 - 6x + 4) \right] dx \\ &= \left[4 \cdot \frac{4}{\pi} \sin\left(\frac{\pi}{4}x\right) - \left(\frac{2x^3}{3} - 3x^2 + 4x\right) \right]_0^2 \\ &= \frac{16}{\pi} - \left(\frac{16}{3} - 12 + 8\right) = \frac{16}{\pi} - \frac{4}{3} \end{aligned}$$

4 : $\begin{cases} 1 : \text{integrand} \\ 2 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \text{(b) Volume} &= \pi \int_0^2 [(4 - f(x))^2 - (4 - g(x))^2] dx \\ &= \pi \int_0^2 \left[(4 - (2x^2 - 6x + 4))^2 - (4 - 4\cos\left(\frac{\pi}{4}x\right))^2 \right] dx \end{aligned}$$

3 : $\begin{cases} 2 : \text{integrand} \\ 1 : \text{limits and constant} \end{cases}$

$$\begin{aligned} \text{(c) Volume} &= \int_0^2 [g(x) - f(x)]^2 dx \\ &= \int_0^2 \left[4\cos\left(\frac{\pi}{4}x\right) - (2x^2 - 6x + 4) \right]^2 dx \end{aligned}$$

2 : $\begin{cases} 1 : \text{integrand} \\ 1 : \text{limits and constant} \end{cases}$