


AP REVIEW SESSION 4

Applications of Derivatives

- Using derivatives to find when a function is Increasing/Decreasing
 - Critical Values
- Relative Extrema/1st Derivative Test
 - Points of Inflection
 - Concavity
 - 2nd Derivative Test
- MVT/Rolle's Theorem
 - EVT
- Tangent Line Approximation
 - Optimization
 - Particle Motion
- Connecting f , f' , and f''
- Sketching f based on f' and f''

No-Calc


2003




7. The graph of f' , the derivative of the function f , is shown above. Which of the following statements is true about f ?
- (A) f is decreasing for $-1 \leq x \leq 1$.
 - (B) f is increasing for $-2 \leq x \leq 0$.
 - (C) f is increasing for $1 \leq x \leq 2$.
 - (D) f has a local minimum at $x = 0$.
 - (E) f is not differentiable at $x = -1$ and $x = 1$.

10. The function f has the property that $f(x)$, $f'(x)$, and $f''(x)$ are negative for all real values x . Which of the following could be the graph of f ?


(A)




(B)




(C)



(D)



(E)



15. Let f be the function with derivative given by $f'(x) = x^2 - \frac{2}{x}$. On which of the following intervals is f decreasing?

(A) $(-\infty, -1]$ only

(B) $(-\infty, 0)$

(C) $[-1, 0)$ only


(D) $(0, \sqrt[3]{2}]$

(E) $[\sqrt[3]{2}, \infty)$

17. Let f be the function given by $f(x) = 2xe^x$. The graph of f is concave down when
- (A) $x < -2$ (B) $x > -2$ (C) $x < -1$ (D) $x > -1$ (E) $x < 0$

x	-4	-3	-2	-1	0	1	2	3	4
$g'(x)$	2	3	0	-3	-2	-1	0	3	2

18. The derivative g' of a function g is continuous and has exactly two zeros. Selected values of g' are given in the table above. If the domain of g is the set of all real numbers, then g is decreasing on which of the following intervals?
- (A) $-2 \leq x \leq 2$ only
 (B) $-1 \leq x \leq 1$ only
 (C) $x \geq -2$
 (D) $x \geq 2$ only
 (E) $x \leq -2$ or $x \geq 2$








21. The second derivative of the function f is given by $f''(x) = x(x - a)(x - b)^2$. The graph of f'' is shown above. For what values of x does the graph of f have a point of inflection?
- (A) 0 and a only (B) 0 and m only (C) b and j only (D) 0, a , and b (E) b , j , and k

28. Let g be a twice-differentiable function with $g'(x) > 0$ and $g''(x) > 0$ for all real numbers x , such that $g(4) = 12$ and $g(5) = 18$. Of the following, which is a possible value for $g(6)$?
- (A) 15 (B) 18 (C) 21 (D) 24 (E) 27

2008

5. The function f given by $f(x) = 2x^3 - 3x^2 - 12x$ has a relative minimum at $x =$
- (A) -1 (B) 0 (C) 2 (D) $\frac{3 - \sqrt{105}}{4}$ (E) $\frac{3 + \sqrt{105}}{4}$

16. If $f'(x) = |x - 2|$, which of the following could be the graph of $y = f(x)$?



19. Let f be the function given by $f(x) = x^3 - 6x^2$. The graph of f is concave up when

- (A) $x > 2$
- (B) $x < 2$
- (C) $0 < x < 4$
- (D) $x < 0$ or $x > 4$ only
- (E) $x > 6$ only

22. If $f'(x) = (x - 2)(x - 3)^2(x - 4)^3$, then f has which of the following relative extrema?

- I. A relative maximum at $x = 2$
- II. A relative minimum at $x = 3$
- III. A relative maximum at $x = 4$

- (A) I only
- (B) III only
- (C) I and III only
- (D) II and III only
- (E) I, II, and III

26. For $x > 0$, f is a function such that $f'(x) = \frac{\ln x}{x}$ and $f''(x) = \frac{1 - \ln x}{x^2}$. Which of the following is true?

- (A) f is decreasing for $x > 1$, and the graph of f is concave down for $x > e$.
- (B) f is decreasing for $x > 1$, and the graph of f is concave up for $x > e$.
- (C) f is increasing for $x > 1$, and the graph of f is concave down for $x > e$.
- (D) f is increasing for $x > 1$, and the graph of f is concave up for $x > e$.
- (E) f is increasing for $0 < x < e$, and the graph of f is concave down for $0 < x < e^{3/2}$.

Calculator Allowed

2003

80. The function f is continuous for $-2 \leq x \leq 1$ and differentiable for $-2 < x < 1$. If $f(-2) = -5$ and $f(1) = 4$, which of the following statements could be false?
- (A) There exists c , where $-2 < c < 1$, such that $f(c) = 0$.
(B) There exists c , where $-2 < c < 1$, such that $f'(c) = 0$.
(C) There exists c , where $-2 < c < 1$, such that $f(c) = 3$.
(D) There exists c , where $-2 < c < 1$, such that $f'(c) = 3$.
(E) There exists c , where $-2 \leq c \leq 1$, such that $f(c) \geq f(x)$ for all x on the closed interval $-2 \leq x \leq 1$.

81. Let f be the function with derivative given by $f'(x) = \sin(x^2 + 1)$. How many relative extrema does f have on the interval $2 < x < 4$?
- (A) One (B) Two (C) Three (D) Four (E) Five

87. The function f has first derivative given by $f'(x) = \frac{\sqrt{x}}{1+x+x^3}$. What is the x -coordinate of the inflection point of the graph of f ?
- (A) 1.008 (B) 0.473 (C) 0 (D) -0.278 (E) The graph of f has no inflection point.

90. For all x in the closed interval $[2, 5]$, the function f has a positive first derivative and a negative second derivative. Which of the following could be a table of values for f ?

(A)

x	$f(x)$
2	7
3	9
4	12
5	16

(B)

x	$f(x)$
2	7
3	11
4	14
5	16

(C)

x	$f(x)$
2	16
3	12
4	9
5	7

(D)

x	$f(x)$
2	16
3	14
4	11
5	7

(E)

x	$f(x)$
2	16
3	13
4	10
5	7

2008

81. The first derivative of the function f is given by $f'(x) = x - 4e^{-\sin(2x)}$. How many points of inflection does the graph of f have on the interval $0 < x < 2\pi$?

(A) Three (B) Four (C) Five (D) Six (E) Seven

82. If f is a continuous function on the closed interval $[a, b]$, which of the following must be true?

(A) There is a number c in the open interval (a, b) such that $f(c) = 0$.

(B) There is a number c in the open interval (a, b) such that $f(a) < f(c) < f(b)$.

(C) There is a number c in the closed interval $[a, b]$ such that $f(c) \geq f(x)$ for all x in $[a, b]$.


(D) There is a number c in the open interval (a, b) such that $f'(c) = 0$.

(E) There is a number c in the open interval (a, b) such that $f'(c) = \frac{f(b) - f(a)}{b - a}$.

x	2.5	2.8	3.0	3.1
$f(x)$	31.25	39.20	45	48.05

83. The function f is differentiable and has values as shown in the table above. Both f and f' are strictly increasing on the interval $0 \leq x \leq 5$. Which of the following could be the value of $f'(3)$?

- (A) 20 (B) 27.5 (C) 29 (D) 30 (E) 30.5



84. The graph of f' , the derivative of the function f , is shown above. On which of the following intervals is f decreasing?


- (A) $[2, 4]$ only
 (B) $[3, 5]$ only
 (C) $[0, 1]$ and $[3, 5]$
 (D) $[2, 4]$ and $[6, 7]$
 (E) $[0, 2]$ and $[4, 6]$

x	3	4	5	6	7
$f(x)$	20	17	12	16	20

86. The function f is continuous and differentiable on the closed interval $[3, 7]$. The table above gives selected values of f on this interval. Which of the following statements must be true?

- I. The minimum value of f on $[3, 7]$ is 12.
- II. There exists c , for $3 < c < 7$, such that $f'(c) = 0$.
- III. $f'(x) > 0$ for $5 < x < 7$.

- (A) I only
- (B) II only
- (C) III only
- (D) I and III only
- (E) I, II, and III



Graph of f'

87. The figure above shows the graph of f' , the derivative of the function f , on the open interval $-7 < x < 7$. If f' has four zeros on $-7 < x < 7$, how many relative maxima does f have on $-7 < x < 7$?

- (A) One
- (B) Two
- (C) Three
- (D) Four
- (E) Five

92. Let f be the function with first derivative defined by $f'(x) = \sin(x^3)$ for $0 \leq x \leq 2$. At what value of x does f attain its maximum value on the closed interval $0 \leq x \leq 2$?

- (A) 0
- (B) 1.162
- (C) 1.465
- (D) 1.845
- (E) 2

FRQ – No Calculator

6. Let $g(x) = xe^{-x} + be^{-x}$, where b is a positive constant.

(a) Find $\lim_{x \rightarrow \infty} g(x)$.

(b) For what positive value of b does g have an absolute maximum at $x = \frac{2}{3}$? Justify your answer.

(c) Find all values of b , if any, for which the graph of g has a point of inflection on the interval $0 < x < \infty$. Justify your answer.